



## Digital Receipt

This receipt acknowledges that Turnitin received your paper. Below you will find the receipt information regarding your submission.

The first page of your submissions is displayed below.

Submission author: Yosep Mau  
Assignment title: Agroteknologi  
Submission title: Efficacy of indigenous *Trichoderma* isolates of West Timor, I...  
File name: III-A2.pdf  
File size: 1.19M  
Page count: 10  
Word count: 7,871  
Character count: 39,700  
Submission date: 11-Apr-2023 06:58PM (UTC+0700)  
Submission ID: 2061485085

Mau et al.  
Egyptian Journal of Biological Pest Control (2023) 32:62  
<https://doi.org/10.1186/s4198-022-05559-x>

Egyptian Journal of  
Biological Pest Control

RESEARCH Open Access

### Efficacy of indigenous *Trichoderma* isolates of West Timor, Indonesia, as biocontrol agents of brown spot (*Drechslera oryzae*) on two upland rice varieties

Yosep Seran Mau<sup>1\*</sup>, Rikardus Suhendra Prayetno<sup>1</sup>, Habil Kaka<sup>1</sup>, Kiki Delfiyanti Naat<sup>1</sup>, Julinda Bendalina Dengga Henuk<sup>1</sup>, Mayavira Veronica Hahuly<sup>1</sup>, Sri Widnugrahani<sup>1</sup> and Yosefma Rovina Yuliaty Ganduk<sup>1</sup>

**Abstract**  
**Background:** Brown spot disease is one of the most destructive foliar diseases of rice. Biological control of this disease is considered more environmentally safe, but the biological control agents have limited availability. The fungus *Trichoderma* can be used as a biocontrol agent for various plant diseases, including brown spots. Indigenous *Trichoderma* species from West Timor were evaluated in this study to reveal their effectiveness in reducing brown spot disease of upland rice. The study was conducted under a screen house condition using a factorial treatment design laid out in a completely randomized design. The two factors were the brown spot disease control method (*T. viride*, *T. harzianum*, *T. hamatum*, fungicide Trivia 73 WP, and a control treatment) and upland rice genotype (Inpage 7 and Breun Senaran 2).  
**Results:** The results showed that *Trichoderma* spp. and the fungicide Trivia 73 WP caused significantly lower AUDPC and apparent infection rate than the control treatment. The total phenol content and agronomic characters (number of reproductive tillers per plant, number of filled grains per panicle, grain yield per plant) were higher in the *Trichoderma* and fungicide treatments than the control treatment. Meanwhile, tannin and saponin levels were non-significantly different among treatments. *Trichoderma* and the fungicide Trivia treatments effectively suppressed brown spot disease of upland rice, and the efficacy was dependent on the rice variety, with a mean efficacy range over the two varieties of 27.05–36.42%. The *Trichoderma* and the fungicide Trivia treatments increased the grain yield per plant differentially in the 2 rice varieties, with a mean grain yield increase over the 2 varieties of 13.0% (*T. viride*), 23.5% (*T. harzianum*), 35.0% (*T. hamatum*), and 19.9% (Trivia 73 WP).  
**Conclusion:** Efficacy of local *Trichoderma* of West Timor and the fungicide Trivia treatments were varietal dependent with *T. hamatum* being the most effective in reducing the brown spot disease and producing the highest grain yield increase over the 2 rice varieties. The local *Trichoderma* species of West Timor have a future potential as biocontrol agents of upland rice brown spot disease, but it is necessary to further evaluate their effectiveness against the disease in replicated field trials.

\*Correspondence: [yosepmau@stafunda.ac.id](mailto:yosepmau@stafunda.ac.id); [yosepmau@yahoo.com](mailto:yosepmau@yahoo.com)  
<sup>1</sup>Agrotechnology Department, Faculty of Agriculture, Universitas Nusa Cendana, Jl. Adisucipto, Wefusa, Kupang 997182001, Indonesia  
Full list of author information is available at the end of the article

Springer Open

© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

# Efficacy of indigenous Trichoderma isolates of West Timor, Indonesia, as biocontrol agents of brown spot (Drechslera oryzae) on two upland rice varieties

*by* Yosep Mau

---

**Submission date:** 11-Apr-2023 06:58PM (UTC+0700)

**Submission ID:** 2061485085

**File name:** III-A2.pdf (1.19M)

**Word count:** 7871

**Character count:** 39700

RESEARCH

Open Access

# Efficacy of indigenous *Trichoderma* isolates of West Timor, Indonesia, as biocontrol agents of brown spot (*Drechslera oryzae*) on two upland rice varieties



Yosep Seran Mau<sup>1,2\*</sup>, Rikardus Suhendra Prayetno<sup>1</sup>, Habil Kaka<sup>1</sup>, Kiki Delfiyanti Naat<sup>1</sup>, Julinda Bendalina Dengga Henuk<sup>1</sup>, Mayavira Veronica Hahuly<sup>1</sup>, Sri Widinugraheni<sup>1</sup> and Yosefina Rovina Yuliaty Gandut<sup>1</sup>

## Abstract

**Background:** Brown spot disease is one of the most destructive foliar diseases of rice. Biological control of this disease is considered more environmentally safe, but the biological control agents have limited availability. The fungus *Trichoderma* can be used as a biocontrol agent for various plant diseases, including brown spots. Indigenous *Trichoderma* species from West Timor were evaluated in this study to reveal their effectiveness in reducing brown spot disease of upland rice. The study was conducted under a screen house condition using a factorial treatment design laid out in a completely randomized design. The two factors were the brown spot disease control method (*T. viride*, *T. harzianum*, *T. hamatum*, fungicide Trivia 73 WP, and a control treatment) and upland rice genotype (Inpago 7 and Breun Senaren 2).

**Results:** The results showed that *Trichoderma* spp. and the fungicide Trivia 73 WP caused significantly lower DPC and apparent infection rate than the control treatment. The total phenol content and agronomic characters (number of reproductive tillers per plant, number of filled grains per panicle, grain yield per plant) were higher in the *Trichoderma* and fungicide treatments than the control treatment. Meanwhile, tannin and saponin levels were non-significantly different among treatments. *Trichoderma* and the fungicide Trivia treatments effectively suppressed brown spot disease of upland rice, and the efficacy was dependent on the rice variety, with a mean efficacy range over the two varieties of 27.05–36.42%. The *Trichoderma* and the fungicide Trivia treatments increased the grain yield per plant differentially in the 2 rice varieties, with a mean grain yield increase over the 2 varieties of 13.0% (*T. viride*), 23.5% (*T. harzianum*), 35.0% (*T. hamatum*), and 19.9% (Trivia 73 WP).

**Conclusion:** Efficacy of local *Trichoderma* of West Timor and the fungicide Trivia treatments were varietal dependent with *T. hamatum* being the most effective in reducing the brown spot disease and producing the highest grain yield increase over the 2 rice varieties. The local *Trichoderma* species of West Timor have a future potential as biocontrol agents of upland rice brown spot disease, but it is necessary to further evaluate their effectiveness against the disease in replicated field trials.

\*Correspondence: yosepmau@staf.un.dana.ac.id; yosepmau@yahoo.com

<sup>1</sup> Agrotechnology Department, Faculty of Agriculture, Universitas Nusa Dana, Jl. Adisucipto, Penfui, Kupang, NTT 85001, Indonesia  
Full list of author information is available at the end of the article

**Keywords:** Rice disease, Brown spot disease, *Trichoderma*, Local species, Biological control, Effectiveness

## Background

Leaf brown spot caused by the fungus *Drechslera oryzae* (Breda de Haan), Subram & Jain [syn. *Helminthosporium oryzae*, *Bipolaris oryzae* (Breda de Haan) Shoemaker], is one of the primary diseases of upland rice that causes high yield losses (IRRI 2009).

In East Nusa Tenggara (ENT) Province, Indonesia, leaf brown spot has long been associated with upland rice farming. The traditional cultivation practices and the poor dry land soil nutrition are the factors that cause the disease to occur more frequently. Consequently, most of the local upland rice cultivars in the region are susceptible and moderately susceptible to leaf brown spot disease, and the yield loss caused by the disease ranged from 10.46 to 56.15% (Mau et al. 2020). Leaf brown spot is, therefore, considered one of the factors contributing to the low upland rice productivity (2 t/ha) in ENT Province, while under controlled and optimum field conditions, the local cultivars can produce a grain yield of 5–6 t/ha (Ndiwa and Mau 2019). Therefore, seeking effective control measures for leaf brown spot disease is necessary to prevent high yield losses in upland rice and hence increase rice productivity.

Brown spot disease can be controlled using various methods (Asghar et al. 2019), including chemical spray, disease-free seeds, use of resistant varieties, biological control, etc. Resistant varieties are the most economical and environmentally sound control measure (Magar 2015). However, the brown spot resistant varieties are limited, and the resistance is frequently short-lived due to the frequent emergence of new virulent races (Arshad et al. 2008). Thus, an integrated control strategy that involves a more environmentally friendly method, such as biological control agents, also needs to be implemented to tackle the disease effectively. For example, the fungus *Trichoderma* sp. is a biological control agent proven to control various plant diseases effectively.

*Trichoderma* is a highly active fungus in the rhizosphere, soil, and the plant canopy environment, and it can also be a parasite to other fungi (Harman et al. 2004). *Trichoderma* produces several types of extracellular proteins and compounds that are fungi toxic and fungi-static to several fungal pathogens and also can stimulate plant growth, so this fungus has received serious attention as a biological control agent (Puyam 2016).

The ability of *Trichoderma* to suppress pathogen and disease development occurs through various mechanisms such as competition (Khalili et al. 2012), mycoparasitism (Dix and Webster 1995), antibiosis (Khalili et al. 2012)

and also by inducing the host plant's resistance (Levy et al. 2015) and promoting the host plant growth (Hassan et al. 2017). Induction of host plant resistance can occur through the production of secondary metabolites such as phenol, saponin, and tannin (Yanti et al. 2018), while plant growth promotion may occur through the production of plant growth hormones, etc. (Alison and Robert 2014).

In an in vitro preliminary study, that several indigenous *Trichoderma* species of West Timor were found inhibiting the aerial growth of the pathogenic fungus *D. oryzae* with an efficacy of 35–50% (Mau et al. 2021). The results indicated the potential of local *Trichoderma* isolates as biological control agents of rice brown spot disease in rice. Evaluating these local *Trichoderma* isolates along with a fungicide spray on different upland rice varieties may lead to finding more effective ways of controlling brown spot disease in upland rice. The objectives of this study were to elucidate the effectiveness of indigenous *Trichoderma* species of West Timor against the brown spot disease on two upland rice genotypes and observe their effects on the agronomic characters on the tested upland rice genotypes.

## Methods

### Study locations and materials

The study was conducted at the Plant Disease Laboratory, Faculty of Agriculture, Universitas Nusa Cendana, and in a screen house in Lasiana Village, Kupang District, East Nusa Tenggara Province, Indonesia. The study was conducted from April to November 2021. Research materials used in the study included *T. viride*, *T. harzianum*, *T. hamatum* (obtained from a collection of Plant Disease Laboratory, Universitas Nusa Cendana), *D. oryzae*, the fungicide Trivia 73 WP, and 2 Indonesian upland rice varieties, i.e., Inpago 7 and Breun Senaren 2.

### Experimental design

The experiment employed a completely randomized design consisting of a factorial treatment design of 2 factors, i.e., upland rice genotype (G) and disease control method (C). The upland rice genotype consisted of 2 varieties (Inpago 7 and Breun Senaren 2), while the disease control method consisted of 5 treatments, i.e., *T. viride*, *T. harzianum*, *T. hamatum*, the fungicide Trivia 73 WP (active ingredients: Fluopicolide 6% and Propineb 66.7%), and a control treatment (without disease control). A total of 15 treatment combinations were evaluated, consisting of 3 replicates.

## Procedures

### Plant cultivation

Rice plants were grown in polybags of 40 cm × 30 cm (height × width) containing soil taken from the surrounding uncultivated field. The soil was taken from 1–20 cm depth, cleaned from the debris, and air-dried for 3 days before being finely ground and filtered to obtain a uniform soil particle. Next, 10 kg air-dried filtered soil was placed into a polybag and then watered to a field capacity level. Four upland rice seeds were planted in each polybag, and only one plant was retained from 2 weeks of age until harvest. A standard upland rice agronomic practice was applied throughout the plant growing cycle. All the plants were maintained in a screen house made from a transparent para net to allow maximum sunlight intensity.

### Preparation of *Trichoderma* solution and foliar application of the treatment

Before being used in the experiment, the *Trichoderma* culture was purified by a single spore method on a potato dextrose agar (PDA) medium for a week at room temperature. Then, the *Trichoderma* was re-identified to confirm their identity by comparing their colony appearance and microscopic characters with the manual description provided by Kubicek dan Harman (2002) and Watanabe (2002). The treatments were applied to the rice plants in a liquid formulation of *Trichoderma* and fungicide through foliar spraying. The *Trichoderma* isolate was grown on a potato dextrose agar (PDA) growth medium for 7 days. Then, 5 disks ( $\theta = 0.6$  cm) of the *Trichoderma* were taken and placed into a 250-ml flask containing 100 ml PD broth medium. The flasks containing the PD broth culture were then continuously shaken at room temperature for 7 days at 150 rpm. The colonized PD broth medium was filtered using a 0.45  $\mu$ m mesh membrane paper. The filtered filtrate was adjusted to a final concentration of  $10^8$  conidia/mL and used as a working solution. The fungicide Trivia 73 WP working solution was prepared by dissolving one gram of the fungicide powder into 1 L distilled water and shaken thoroughly to obtain a homogeneous working solution with a final concentration of 1 g/L.

*Trichoderma* and the fungicide Trivia 73 WP were administrated by foliar spray 19 days after planting (2 days before plant inoculation using *D. oryzae*). The working solutions of *Trichoderma* and fungicide Trivia were sprayed onto the rice plant's leaves until runoff occurred. The foliar application was carried out at around 8 to 9 am. The treatment applications were conducted 4 times on a two-weekly basis. Plants that received no *Trichoderma* and fungicide sprays served as the control (no disease control treatment).

### Preparation of *D. oryzae* suspension and plant inoculation

The fungus *D. oryzae* was obtained from infected rice fields. The diseased leaf disks were cultured on a PDA medium for 7 days, and a single spore was isolated and further grown on a PDA medium for another 7 days under room temperature for purification and identification of *D. oryzae*. Identification was done by comparing the colony appearance and microscopic characters with the authentic description provided by Ellis (1971) and Agarwal et al. (1989). A PDA medium was used to propagate the *D. oryzae* isolate for plant inoculation. After 14 days, the fungus mycelia mats were gently scraped with a spatula and added with 10 ml sterile water containing 0.02% Tween 20 to obtain the conidial suspension. The conidial concentration of the stock solution was determined using a hemocytometer and then adjusted to  $1 \times 10^6$  conidia/ml for plant inoculation. Plant inoculation was conducted 21 days after planting (2 days after foliar sprays of *Trichoderma* sp. and the fungicide). All tested plants were inoculated by foliar spraying until runoff occurred. Inoculation was done from 8 to 9 am.

### Observation

Observation was done on individual plants, including disease severity, phenol content, tannin and saponin levels, the number of reproductive tillers per plant, the number of filled grains per panicle, and grain yield per plant. Disease severity was assessed every week from 7 days after inoculation until the early flowering stage. The disease was assessed based on a disease scale according to the standard evaluation system of IRRI (2013) and was then used to calculate the disease severity level using the formula:

$$DS = \frac{\sum (n \times v)}{Z \times N} \times 100\%$$

where DS = disease severity,  $n$  = number of leaves with a certain disease score,  $v$  = disease score,  $Z$  = the highest observed disease score, and  $N$  = the total number of leaves observed in each plant. The mean disease severity of each plant was averaged from all leaves in the plant.

A disease severity recordings were used to calculate the area under the disease progress curve (AUDPC) based on the formula of Campbell and Madden (1990).

$$AUDPC = \sum_i^{n-1} \frac{Y_i + Y_{i+1}}{2} (t_{i+1} - t_i)$$

where  $Y_i$  = disease severity at the  $i$ th observation,  $n$  = the last disease assessment (number of assessment), and  $t_i$  = time period of assessment at the  $i$ th observation.

**Table 1** Mean square values of epidemiological, phenol content, yield, and yield-attributing parameters of two rice genotypes treated with different brown spot disease control methods

Source of variation	df	Epidemiological parameters		Total phenol content	Yield and yield-contributing parameter		
		AUDPC	rt		NRTP	NFGP	GYP
Genotype (G)	1	208,484.98**	0.00021*	0.14784**	607,500**	1128.53**	54.68**
Control method (C)	4	224,706.62**	0.00006**	0.12559**	11.283**	579.46**	27.75**
G*C	10	4024.40.ns	0.00042*	0.52327**	1.250.ns	366.24**	22.05**
Error	20	13,046.33	0.00004	0.00012	1.667	49.02	5.55
CV (%)	–	11.31	19.73	0.60	6.47	6.01	11.93

df = degree of freedom, AUDPC = area under the disease progress curve, rt = apparent infection rate, NRTP = number of reproductive tiller per plant, NFGP = number of filled grain per panicle, GYP = grain yield per plant, CV = coefficient of variation, \*\*highly significant ( $P < 0.01$ ), \*Significant ( $P < 0.05$ ), ns not significant ( $P > 0.05$ )

The mean apparent infection rate was calculated based on the time-varying apparent rate (Madden et al. 2007) as follows:

$$r_{t+1} = \left[ \ln \left( \frac{y_{t+1}}{1 - y_{t+1}} \right) - \ln \left( \frac{y_t}{1 - y_t} \right) \right] / (t_{t+1} - t_t)$$

where  $y_t$  = disease severity at the  $t$ th observation, and  $t_t$  = time period of assessment at the  $t$ th observation.

Total phenol content and tannin and saponin levels were determined on leaves of the tested rice plants 3 days before the final disease assessment. Five grams of fresh leaf sample of each plant was taken and used for the analysis in the Laboratory of Biosains, Universitas Nusa Cendana. Total phenol content was determined based on the method of McDonald et al. (2001), while saponin and tannin levels were qualitatively assessed based on the method of Syahadat and Azis (2012).

Agronomic data were recorded at harvest. The number of reproductive tillers was obtained by counting the reproductive tiller per plant, while the number of grains per panicle was averaged from the number of grains of all panicles within a plant. The grain yield was measured as grain weight ( $\pm 14\%$  moisture content) per plant.

#### Data analysis

Quantitative data such as AUDPC, apparent infection rate, total phenol content, and agronomical data were subjected to ANOVA followed by an LSD post hoc test, while saponin and tannin levels were descriptively analyzed. Correlation analysis was performed to see the associations among variables. The ANOVA was performed using GenStat 12th (VSN International 2009), while the correlation analysis was conducted using PAST ver. 3 (Hammer et al. 2001).

### Results

#### Brown disease severity and efficacy of the treatments

Ten recordings of brown spot disease severity were made during the study. The disease symptoms started to appear

from the first week after inoculation. At the last assessment, the disease severity ranged on the 2 tested varieties which were 19.1–26.6 and 19.6–26.1%, respectively.

Data in Table 1 summarize the results of factorial ANOVA of epidemiological and agronomical parameters. The single factor genotype and disease control method significantly or highly significantly affected the AUDPC and apparent infection rate. In contrast, the interaction between the 2 factors caused non-significant effect on AUDPC but significantly affected the apparent infection rate.

Data (\*) in Table 2 show significant AUDPC among treatments within every single factor of rice genotype (G) and disease control method (C). The overall mean AUDPC over the 2 varieties (mean G) was highest in the control treatment (1345.03%.day), while the lowest was in *T. hamatum* (855.20%.day), but did not differ significantly from that of *T. harzianum* (908.59%.day). Consistent with the mean genotype, the highest AUDPC within each genotype was observed on the control treatment, and the lowest was on *T. hamatum* and *T. harzianum* applications. The overall mean AUDPC of the disease control method factor was higher in Breun Senaren 2 (1093.41%.day) than Inpago 7 (926.68%.day). Within each disease control method, Inpago 7 consistently showed lower AUDPC than Breun Senaren 2, indicating that the former had a higher brown spot resistance than the latter.

Apparent infection rates also varied significantly among treatments. The treatment combination of Breun Senaren 2  $\times$  *T. viride* showed the lowest apparent infection rate (0.0077 disease/unit/day), while the highest was on Inpago 7  $\times$  control treatment (0.0343 disease/unit/day). On average, the apparent infection rate in Inpago 7 (0.019 disease/unit/day) was higher than Breun Senaren 2 (0.016 disease/unit/day). The mean apparent infection rate of the disease control method over the 2 rice genotypes was lowest in *T. harzianum* (0.0093 disease/unit/day), and the highest was on the control treatment (0.0297 disease/unit/day).

**Table 2** Means of AUDPC, apparent infection rate, and efficacy of leaf brown spot disease under combined treatments of disease control method and upland rice genotype

Control method (C)	AUDPC (%.day)			Apparent infection rate (disease/unit/day)			Efficacy (%)		
	Rice genotype (G)			Rice genotype (G)			Rice genotype (G)		
	Inpago 7	Breun Senaren 2	Mean (G)	Inpago 7	Breun Senaren 2	Mean (G)	Inpago 7	Breun Senaren 2	Mean (G)
<i>T. viride</i>	866.83 a A	1095.69 b B	981.26 b	0.0126 a A	0.0077 a B	0.0101 bc	31.71 a B	22.88 a A	27.05 a
<i>T. harzianum</i>	806.28 a A	1010.91 ab B	908.59 ab	0.0103 a A	0.0084 a A	0.0093 a	36.48 a B	28.85 a A	32.45 a
<i>T. hamatum</i>	807.21 a A	903.19 a B	855.20 a	0.0171 b A	0.0207 b A	0.0189 c	36.40 a A	36.43 b A	36.42 b
Trivia 73 WP	883.83 a A	1036.46 ab B	960.15 b	0.0209 b A	0.0181 b A	0.0195 c	30.37 a A	27.05 a A	28.62 a
No control	1269.26 b A	1420.80 c B	1345.03 c	0.0343 c A	0.0252 c B	0.0297 d			
Mean (C)	926.68 A	1093.41 B		0.0190 A	0.0160 B				

<sup>1</sup> Lowercase denotes comparison within the same column, and uppercase indicates comparison within the same row. Values within the same column/row with the identical lowercase or uppercase are not significantly different based on the LSD (0.05) post hoc test

\*Determined as the percentage of decrease in AUDPC from that of the control treatment

Data in Table 2 also show that the disease control method using *Trichoderma* sp. and fungicide sprays effectively reduced the disease AUDPC than the control treatment. The efficacy of the *Trichoderma* and the fungicide Trivia varied significantly between the 2 rice varieties. Over the 2 varieties, the highest efficacy (36.42%) was obtained from *T. hamatum* spray. In comparison, the effectiveness of the other 3 treatments was 27.05% (*T. viride*), 28.62% (fungicide Trivia 73 WP), and 32.45% (*T. harzianum*), which did not significantly differ from each other. Efficacy of *Trichoderma* and the fungicide Trivia was statistically similar on Inpago 7 but was significantly different on Breun Senaren 2 with *T. hamatum* the most effective.

#### Phenol, tannin, and saponin levels

Total phenol content was highly significantly affected by the interaction of disease control method and rice genotype (Table 1). Similarly, the single factor treatments showed a highly significant effect on the total phenol content. Data in Table 3 show that the treatment combination of *T. hamatum* × Inpago 7 produced the highest phenol content (2.40 mg GAE/g), followed by *T. harzianum* × Inpago 7 (2.07 mg GAE/g) and the fungicide Trivia 73 WP × Breun Senaren 2 (2.00 mg GAE/g) in the second place. Meanwhile, the lowest phenol content was recorded on the control × Inpago 7 treatment combination (1.42 mg GAE/g). On the overall mean of the 2 varieties, *T. harzianum* spray produced the highest phenol

<sup>17</sup> content (2.01 mg GAE/g), which was significantly higher than other treatments, while the lowest was observed on the control treatment (1.61 mg GAE/g). The mean phenol content of the the five disease control treatments over the two varieties differed significantly from each other. The overall mean phenol content of disease control method treatment was higher in Inpago 7 (1.91 mg GAE/g) than Breun Senaren 2 (1.76 mg GAE/g). In contrast to total phenol content, tannin was equally produced in all treatment combinations at a high level in both varieties (Table 3). Meanwhile, saponin was produced in low to medium levels in both varieties. The control treatment had a low level of saponin in both varieties, which was also similar to the treatment combinations of *T. hamatum* × Inpago 7, *T. viride* × Breun Senaren 2, and *T. harzianum* × Breun Senaren 2, while the rest treatment combinations produced medium levels of saponin.

#### Number of reproductive tiller per plant and number of filled grain per panicle

<sup>4</sup> Interaction of rice variety (G) and disease control method (C) caused non-significant effect on the number of reproductive tillers per plant. In contrast, both the single factor treatments caused a highly significant effect on the trait (Table 1). Mean reproductive tillers per plant over the 2 genotypes were highest in the fungicide Trivia 73 WP (21.33 tillers), but it was non-significantly different from *Trichoderma* sp. applications (Table 4). In comparison,

**Table 3** Means of leaf phenol, tannin, and saponin levels under combined treatments of disease control method and upland rice genotype

Control method (C)	Total phenol (mg GAE/g)						Tannin level*		Saponin level*	
	Rice genotype (G)						Rice genotype		Rice genotype	
	Inpago 7	Breun Senaren 2	Mean (G)		Inpago 7	Breun Senaren 2	Inpago 7	Breun Senaren 2		
<i>T. viride</i>	1.83 A	c B	1.81 b	b	1.82 b	+++	+++	++	+	
<i>T. harzianum</i>	2.07 A	d B	1.94 c	c	2.01 e	+++	+++	++	+	
<i>T. hamatum</i>	2.40 A	e B	1.27 a	a	1.84 c	+++	+++	+	++	
Trivia 73 WP	1.80 A	b B	2.00 d	d	1.90 d	+++	+++	++	++	
No disease control	1.42 A	a B	1.80 b	b	1.61 a	+++	+++	+	+	
Mean (C)**	1.91 A		1.76 B							

<sup>1</sup> Lowercase denotes comparison within the same column, and uppercase denotes comparison within the same row. Values within the same column/row with the same lowercase or uppercase are not significantly different based on the LSD (0.05) post hoc test. \*Qualitatively assessed: ++ low level, +++ medium level, ++++ high level

**Table 4** Means of number of reproductive tillers per plant and number of grain per panicle under combined treatments of disease control method and upland rice variety

Control method (C)	Reproductive tiller per plant			Number of grain per panicle/tiller			Grain yield per plant (g)		
	Rice Genotype (G)			Rice Genotype (G)			Rice Genotype (G)		
	Inpago 7	Breun Senaren 2	Mean (G)	Inpago 7	Breun Senaren 2	Mean (G)	Inpago 7	Breun Senaren 2	Mean (G)
<i>T. viride</i>	23.00 a A	14.00 b B	18.50 a	118.33 b A	121.22 bc A	119.78 c	18.50 a A	19.00 b B	18.75 b
<i>T. harzianum</i>	26.00 b A	16.00 b B	21.00 b	130.44 c A	93.78 a B	112.11 b	25.50 c B	16.33 a A	20.92 bc
<i>T. hamatum</i>	25.33 b A	15.67 ab B	20.50 b	114.78 ab A	112.78 b A	113.78 b	22.50 b A	22.33 c A	22.42 c
Trivia 73 WP	25.67 b A	17.00 bc B	21.33 b	140.33 d A	122.78 c B	131.56 d	21.17 b B	18.67 b A	19.92 b
No disease Control	22.33 a A	14.67 a B	18.50 a	109.44 a A	101.44 a A	105.44 a	17.83 a B	15.67 a A	16.75 a
Mean (C)**	24.47 A	15.47 B		122.67 A	110.40 B		21.10 B	18.40 A	


<sup>1</sup> Lowercase denotes comparison within the same column, and uppercase indicates comparison within the same row. Values within the same column/row with the identical lowercase or uppercase are not significantly different based on the LSD (0.05) post hoc test

the lowest number of reproductive tillers was recorded on the control treatment (18.50 tillers).

On average, the number of reproductive tillers per plant of Inpago 7 (24.47) was higher than Breun Senaren 2 (15.47 tillers). A similar situation did happen within each disease control method (Table 4).

Interaction of disease control method and rice variety highly affected ( $P < 0.01$ ) the number of filled grains per panicle (Table 1). The highest number of grain per panicle was observed on the treatment combination of Trivia 73 WP × Inpago 7 (140.33 grains), followed by *T. harzianum* × Inpago 7 (130.44 grains) and Trivia 73 WP × Breun Senaren 2 (121.22 grains) in the second and





the third places, respectively (Table 4). Meanwhile, the lowest number of grains per panicle was observed on the treatment combination of no disease control × Inpago 7 (109.44 grains) and no disease control × Breun Senaren 2 (101.44 grains) (Table 4).

On the overall mean of the 2 varieties, the fungicide spray using Trivia 73 WP produced the highest number of grains per panicle (131.56 grains), followed by *T. viride* (119.78 grains) in the second, and *T. harzianum* and *T. hamatum* in the third place. On the other hand, the control treatment produced the lowest number of grains per panicle (105.44 grains). On the overall means of disease control method, Inpago 7 produced a higher number of grains per panicle (122.67 grains) than Breun Senaren 2 (110.40 grains) (Table 4).

#### Grain yield per plant and increase (%) of grain yield per plant

The ANOVA results revealed that the interaction between the disease control method and rice genotype significantly affected grain yield per plant (Table 1). Mean grain yield per plant is presented in Table 4.

On the Inpago 7 genotype, *T. harzianum* produced the highest mean grain yield/plant (25.50 g), followed by *T. hamatum* (22.50 g) and Trivia 73 WP (21.17 g). In contrast, the control treatment produced the lowest grain yield/plant (17.83 g). *T. hamatum* had the highest mean grain yield plant/plant on Breun Senaren 2 (22.33 g), followed by *T. viride* (19.0 g), Trivia 73 WP (18.67 g), *T. harzianum* (16.33 g), and the control treatment (15.67 g). On

the overall mean of the two varieties, *T. hamatum* produced the highest grain yield/plant (22.42 g), followed by *T. harzianum* (20.92 g), Trivia 73 WP (19.92 g), and *T. viride* (18.75 g). The control treatment produced the lowest grain/plant (16.75 g).

The results of grain yield per plant indicate that foliar applications of *Trichoderma* species and fungicide Trivia 73 WP could increase the grain yield of tested upland rice varieties compared to no disease control treatment. The percentage of grain yield increase is presented in Fig. 1, where the percentage of grain increase caused by *Trichoderma* and Trivia fungicide applications was dependent on the variety. On Inpago 7, *T. hamatum* caused the highest increase of grain yield (42.8%), while on Breun Senaren 2, *T. hamatum* produced the highest increase of grain yield per plat (42.9%). Over the 2 varieties, the percentage of grain yield increase was highest on *T. hamatum*, while *T. viride*, *T. harzianum*, and Trivia 73 WP caused a statistically similar percentage of grain yield per plant.

#### Correlation analysis

Correlation analysis was performed to see the association among the studied variables, employing the replicated variable's data of the treatment combinations (Table 5). AUDPC had a positive correlation with apparent infection rate ( $r=0.47$ ), and it had a negative correlation with the reproductive tiller ( $r=-0.49$ ), the number of filled grain ( $r=-0.49$ ), and grain yield ( $r=-0.55$ ), but it had non-significant correlation with phenol content

**Table 5** Correlation analysis of the observed variables

	AUDPC	AIR	TPC	NRTTP	NFGP	GYP
AUDPC	1	0.47*	-0.33	-0.49*	-0.46*	-0.55*
AIR		1	-0.49*	-0.09	-0.10	-0.08
TPC			1	0.35	0.13	0.17
NRTTP				1	0.53*	0.49*
NFGP					1	0.54*
GYP						1

AUDPC = Area under disease progress curve, AIR = apparent infection rate, TPC = total phenol content, NRTTP = number of the reproductive tiller per plant, NFGP = number of filled grain per panicle, GYP = grain yield per plant. Values in the table are correlation coefficients. \* = significant based on t test ( $P < 0.05$ )

( $r = -0.33$ ). The apparent infection rate was negatively correlated with phenol content ( $r = -0.49$ ) and AUDPC, but it did not correlate with other variables. Total phenol had no association with other variables except with apparent infection rate (a negative correlation).

The number of reproductive tillers correlated positively with both the number of filled grain ( $r = 0.53$ ) and grain yield ( $r = 0.49$ ), but it negatively correlated with AUDPC. Grain yield per plant correlated positively with both the number of filled grains per panicle ( $r = 0.49$ ) and the number of filled grain per panicle ( $r = 0.541$ ), but it was negatively correlated with AUDPC (Table 5).

### Discussion

In this study, foliar applications of local *Trichoderma* species and the fungicide Trivia 73 WP were investigated for their effectiveness in suppressing the brown leaf spot in two genotypes of upland rice. The disease control treatments using *Trichoderma* sp. and fungicide treatments were able to suppress the leaf brown spot disease progress on two upland rice varieties as indicated by their lower AUDPC than in the control treatment. In addition, the lower AUDPC in the *Trichoderma* sp. and the fungicide applications was consistent with the lower mean apparent infection rate of these treatments than the control treatment. The efficacy of *Trichoderma* sp. and the fungicide treatments ranged from 27.05 to 36.42%. Foliar application of *T. hamatum* was the most effective against the disease, while *T. viride* and *T. harzianum* were as effective as fungicide spray using Trivia 73 WP. The ability of *Trichoderma* sp. to suppress leaf brown spot in the current study is consistent with previous study results by Khalili et al. (2012). The results also confirmed the effectiveness of *Trichoderma* application against blast diseases of rice (Chou et al. 2020) and various crop diseases (Soesilo et al. 2019).

Results of the present study revealed that total phenol content was higher in plants receiving *Trichoderma* applications than in the control treatment. The results also showed that *Trichoderma* application reduced the

AUDPC and apparent infection rate, supported by the positive correlation between the 2 parameters ( $r = 0.47$ ). Meanwhile, there was a significantly negative correlation between phenol content and the apparent infection rate ( $r = -0.49$ ), indicating the involvement of phenol in suppressing the disease progression rate. Although previous studies had reported phenol involvement in induced resistance against *D. oryzae* (Hassan et al. (2017), the novelty of the present study results is that phenol production of local *Trichoderma* of West Timor and Trivia applications was varietal dependent. This suggests that phenol involvement in suppressing leaf brown spots depends on the rice variety and the *Trichoderma* species, implying different mechanisms of host-pathogen interaction, resulting in various resistance levels.

All treatment combinations produced similarly high-level tannin content, indicating that tannin may not involve in the host plant response against *D. oryzae* and, presumably, the high-level tannin content was produced constitutively, which is in line with Dixon (2001). In contrast, Yanti et al. (2018) reported the involvement of tannin in resistance against dieback disease (*Botryodiplodia theobromae*) of cocoa seedlings. Unlike tannin, saponin was produced in low to medium levels with no apparent pattern among rice genotype and disease control treatments, which renders its involvement in resistance to brown spot disease inconclusive. Nevertheless, previous studies demonstrated that saponin is involved in biochemical resistance against various plant diseases (Yanti et al. 2018).

The number of productive tillers per plant was negatively correlated ( $r = -0.49$ ) with AUDPC, indicating that an increase in AUDPC will be followed by a decrease in productive tiller number and vice versa. Presumably, the increase in AUDPC caused a decline in the plant's photosynthetic capacity since the brown spot symptoms mainly occurred on rice leaves, which reduces the photosynthetic area of the plant. This will, in turn, cause a decline in the number of reproductive tillers (Dariush et al. 2020). The ability of *Trichoderma* and fungicide

applications to produce a higher number of tillers than the control treatment was presumably related to the lower AUDPC, and hence a higher leaf photosynthetic capacity they possessed. The genetic factor may contribute to the significant difference between varieties in the reproductive tiller number as the ability to produce reproductive tiller is varietal dependent (Ndiwa and Mau 2019).

The treatment combinations highly affected the number of filled grains, implying that this trait is highly variable among the disease control treatments and the rice variety. The differences in disease severity may cause different photosynthetic capacities of the rice plants, which resulted in a significant grain number per panicle among treatments. A previous study by Hassan et al. (2017) showed that *Trichoderma* foliar application reduced the brown spot infection and caused an increase in agronomic characters, including the number of discolored (filled) grain.

The grain yield was affected significantly by the disease control method and rice genotype interaction, which may reflect the genetic potential of the tested rice genotypes. The expression of this genetic potential is further influenced by environmental factors, i.e., disease control treatments using *Trichoderma* and the fungicide. The grain yield of the 2 varieties in the *Trichoderma* and fungicide treatments was significantly higher than the control treatment. The lower grain yield in the control treatment may be caused by the decrease in both reproductive tiller number and number of grains per panicle as the direct effect of leaf infection by the disease, which reduced the photosynthetic capacity of rice plants (Pantha et al. 2017). This presumption is supported by the negatively significant correlation between AUDPC with, respectively, grain yield ( $r = -0.55$ ), the number of the reproductive tiller ( $r = -0.49$ ), the number of filled grains per panicle ( $r = -0.46$ ), and the positive correlation between grain yield with, respectively, the number of the reproductive tiller ( $r = 0.49$ ) and the number of filled grains ( $r = 0.54$ ).

A previous study by Hassan et al. (2017) showed that *Trichoderma* foliar application reduced the brown spot infection but increased the agronomic characters (leaf chlorophyll content, plant height, panicle length, the number of filled grain, and grain yield) as compared to the non-treated plants. The ability of *Trichoderma* to improve the agronomic characters is consistent with previous findings that *Trichoderma* can stimulate the host plant growth (Hassan et al. 2017).

The present study results revealed that the efficacy of *Trichoderma* and fungicide treatments against brown spot ranged from 27.05 to 36.42%, while the grain yield of the plants receiving the treatments was

about 13–35% higher than the control. This discrepancy indicated that the grain yield observed in this study may reflect the genetic potential of the tested rice genotypes, which may or may not be directly related to leaf brown spot severity. The higher grain yield in the *Trichoderma*-treated plants than the control might be occurred through both the reduced leaf brown spot infection and the mechanism of plant growth stimulation. These included an increased number of reproductive tillers and filled grain. As a comparison, previous study (Dariush et al. 2020) confirmed that brown spot infection reduced the grain yield by reducing the reproductive tiller number and the number of filled grains. However, the host plant's growth stimulation was not examined physiologically in the present study, which is the limitation of the current study that needs to be considered in future works.

## Conclusions

The indigenous *Trichoderma* species isolated from West Timor and the fungicide Trivia 73 WP effectively reduced brown spot disease of upland rice, and the efficacy was dependent on the rice variety, with a mean efficacy range over the 2 varieties from 27.05 to 36.42%. The *Trichoderma* and the fungicide Trivia treatments increased the grain yield per plant differentially in the 2 rice varieties, with a mean grain yield increase over the 2 varieties of 13.0% (*T. viride*), 23.5% (*T. harzianum*), 35.0% (*T. hamatum*), and 19.9% (Trivia 73 WP). The local *Trichoderma* species of West Timor have a future potential as biocontrol agents of brown spot disease of upland rice as they were able to suppress the disease and increase the grain yield of upland rice under a screen house condition. However, before being further employed as potential biological agents for brown spot disease, future works are needed to evaluate the effectiveness of this local *Trichoderma* against the disease in replicated field trials.

## Abbreviations

ENT: East Nusa Tenggara; WP: Wettable powder; DS: Disease severity; AUDPC: Area under the disease progress curve; ANOVA: Analysis of variance; AIR: Apparent infection rate; TPC: Total phenol content; NRTP: Number of the reproductive tiller per plant; NFGP: Number of filled grain per panicle; GYP: Grain yield per plant; LSD: Least significant difference.

## Acknowledgements

We are very grateful to the Laboratory of Plant Disease, Universitas Nusa Cendana, for providing the *Trichoderma* isolates, the Indonesian Rice Research Institute for providing the upland rice seeds of Inpago 7 variety, and East Flores Department of Agriculture for providing the seeds of Breun Senaren 2 variety. We extend special thanks to Mrs. Rina Sula for her excellent assistance in the laboratory work.

## Author contributions

YSM organized the research team, provided the research supplies, and contributed to all experiment stages. RSP, KDN, and HK contributed to all

stages of the experiment. JBDH and SW assisted in designing the experiment, performed laboratory analysis, and performed data analysis and interpretation. MVH assisted in data analysis and interpretation and assisted in drafting and finalizing the manuscript, and YRYG assisted in designing the experiment and assisted in data analysis and interpretation. All authors read and approved the final manuscript.

#### Funding

This research was funded by Universitas Nusa Cendana through the professorship research scheme year 2021. (Contract number: 178/PP/2021).

5

#### Availability of data and materials

The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

#### Declarations

#### Ethics approval and consent to participate

Not applicable.

#### Consent for publication

The manuscript has not been published completely or in part elsewhere.

#### Competing interests

The authors declare no conflicts of interest.

#### Author details

<sup>1</sup>Agri Technology Department, Faculty of Agriculture, Universitas Nusa Cendana, Jl. Adisucipto, Penfui, Kupang, NTT 85001, Indonesia. <sup>2</sup>Integrated Field Laboratory of Archipelagic Dryland, Universitas Nusa Cendana, Jl. Adisucipto, Penfui, Kupang, NTT 85001, Indonesia.

Received: 18 February 2022 Accepted: 17 May 2022

Published online: 30 May 2022

#### References

- Agarwal PC, Mortensen CN, Mathur SB (1989) Seed-borne diseases and seed health testing of rice. *Phytopathol Pap* 30:106
- Alison S, Robert H (2014) Applications of *Trichoderma* in plant growth promotion. In: *Biotechnology and biology of Trichoderma*, pp 415–428. <https://doi.org/10.1016/B978-0-444-59576-8.00031-X>
- Arshad HMI, Khan JA, Jamil FF (2008) Screening of rice germplasm against blast and brown spot diseases. *Pak J Phytopathol* 20(1):52–57
- Asghar M, Baig MMQ, Chaudhary S, Iqbal MF, Ali MA (2019) Evaluation of difenoconazole along with macronutrients spray for the control of brown leaf spot (*Bipolaris oryzae*) disease in rice (*Oryza sativa*) crop. *Sarhad J Agric* 35(1):1–6. <https://doi.org/10.17582/journal.sja/2019/35.1.1.6>
- Campbell CL, Madden LV (1990) *Introduction to plant disease epidemiology*. Wiley, New York
- Chou C, Castilla N, Hadi B, Tanaka T, Chiba S, Sato I (2020) Rice blast management in Cambodian rice fields using *Trichoderma harzianum* and a resistant variety. *Crop Prot* 135:104864. <https://doi.org/10.1016/j.cpro.2019.104864>
- Dariush S, Darvishnia M, Ebadi AA, Padasht-Dehkaei F, Bazgir E (2020) Screening rice genotypes for brown spot resistance along with yield attributing characters and its association with morphological traits. *J Crop Prot* 9(3):381–393
- Dix NJ, Webster J (1995) *Fungal ecology*. Chapman & Hall
- Dixon RA (2001) Natural products and plant disease resistance. *Nature* 411:843–847
- Ellis MB (1971) *Dematiaceae hyphomycetes*. CMI, Kew, England.
- Hammer Ø, Harper DAT, Ryan PD (2001) PAST: paleontological statistics software package for education and data analysis. [http://palaeo-electronica.org/2001\\_1/past/issue1\\_01.htm](http://palaeo-electronica.org/2001_1/past/issue1_01.htm). Accessed 26 Oct 2019
- Harman GE, Howell CR, Viterbo A, Chet I, Lorito M (2004) *Trichoderma* species—opportunistic, avirulent plant symbionts. *Nat Rev* 2:43–56
- Hassan AA, Kalboush ZA, Saleh MM (2017) Efficacy of *Trichoderma* spp as biocontrol agents against rice brown spot disease and bioc hemical approaches. *J Plant Prot Path Mansoura Univ* 8(12):677–686
- IRRI (2009) Brown spot (rice facts sheet). International Rice Research Institute, Los Banos
- IRRI (2013) Standard evaluation system for rice, 5th edition. International Rice Research Institute, Los Banos
- Khalili E, Sadravi M, Naeimi S (2012) Biological control of rice brown spot with native isolates of three *Trichoderma* species. *Braz J Microbiol* 43(1):297–305. <https://doi.org/10.1590/S1517-838220120001000035>
- Kubicek CP, Harman GE (2002) *Trichoderma* and *Gliocladium*. Vol. 1. Basic Biology, Taxonomy and Genetics. Taylor & Francis
- Levy NO, Harel YM, Haile ZM, Elad Y, Rav-David E, Jurkevitch E, Katan J (2015) Induced resistance to foliar diseases by soil solarization and *Trichoderma harzianum*. *Plant Pathol* 64(2):365–374. <https://doi.org/10.1111/ppa.12255>
- Magar PB (2015) Screening of rice varieties against brown leaf spot disease at Jyotinagar, Chitwan, Nepal. *Int J Appl Sci Biotechnol* 3(1):56–60. <https://doi.org/10.3126/ijasbt.v3i1.12014>
- Mau YS, Ndiwa ASS, Oematan SS (2020) Brown spot disease severity, yield and yield loss relationships in pigmented upland rice from East Nusa Tenggara, Indonesia. *Biodiversitas* 21(4):2625–2634
- Mau YS, Hahuly MV, Henuk JBD, Widinugraheni S (2021) Efficacy of *Trichoderma* species from West Timor against *Dreschlera oryzae* in the laboratory. Research Report. Research and Community Service Institute, Universitas Nusa Cendana, Indonesia (in Indonesian)
- McDonald S, Prenzler PD, Autolovich M, Robards K (2001) Phenolic content and antioxidant activity of olive extracts. *Food Chem* 73:73–84. [https://doi.org/10.1016/S0308-8146\(00\)00288-0](https://doi.org/10.1016/S0308-8146(00)00288-0)
- Ndiwa ASS, Mau YS (2019) Yield and yield component performances of local pigmented upland rice cultivars from East Nusa Tenggara, Indonesia in three locations. *Int J Trop Drylands* 3(2):49–55
- Pantha P, Shrestha SM, Manandhar HK, Gaire SP, Aryal L, Yadav DR (2017) Evaluation of rice genotypes for resistance against brown spot disease caused by *Bipolaris oryzae*. *Int J Curr Res* 9(04):48562–48569
- Puyam A (2016) Advent of *Trichoderma* as a bio-control agent—a review. *J Appl Nat Sci* 8(2):1100–1109. <https://doi.org/10.31018/jans.v8i2.927>
- Soesanto L, Mugiastuti E, Manan A (2019) Raw secondary metabolites application of two *Trichoderma harzianum* isolates towards vascular streak dieback on cocoa seedlings. *Pelita Perkebunan* 35(1):22–32 (in Indonesian)
- Syahadat RM, Aziz SA (2012) Effect of media composition and fertigation with organic fertilizer to bioactive contained of orange jassmine (*Murraya paniculata* (L) Jack) leaf in seedling. *Bull Littro* 23(2):142–147 (in Indonesian)
- VSN International (2009) *GenStat for windows*, 12th edn. VSN International Ltd., Hertfordshire
- Watanabe T (2002) *Pictorial atlas of soil and seed fungi: morphologies of cultured fungi and key to species*, 2nd edn. CRC Press LLC, Boca Raton
- Yanti LA, Achmad A, Khumaida N (2018) Resistance mechanisms of white jaboron seedlings (*Anthocephalus cadamba*) against *Botryodiplodia theobromae* causing dieback disease. *Biodiversitas* 19(4):1441–1459. <https://doi.org/10.13057/biodiv/d190434>

#### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

# Efficacy of indigenous Trichoderma isolates of West Timor, Indonesia, as biocontrol agents of brown spot (*Drechslera oryzae*) on two upland rice varieties

## ORIGINALITY REPORT

13%

SIMILARITY INDEX

%

INTERNET SOURCES

13%

PUBLICATIONS

%

STUDENT PAPERS

## PRIMARY SOURCES

- 1 Yohanis Ngongo, Tony Basuki, Bernard deRosari, Evert Y. Hosang et al. "Local Wisdom of West Timorese Farmers in Land Management", Sustainability, 2022  
Publication 3%
- 2 Amina M. Ibrahim, Mostafa Y. Morad, Salwa A. H. Hamdi, Mona F. Fol. "Biocontrol potential of Chitosan extracted from *Procambarus clarkii* (Crustacea: Cambaridae) against *Eobania vermiculata* snails (Muller 1774) in Egypt", Egyptian Journal of Biological Pest Control, 2022  
Publication 2%
- 3 "Transforming Coastal Zone for Sustainable Food and Income Security", Springer Science and Business Media LLC, 2022  
Publication 1%
- 4 "Association analysis of F2 generation in rice (*Oryza sativa* L.) under salinity condition", 1%

- 5** Eman E. S. El-Sharkawy, Eman Abdelrazik. "Biocontrol of Fusarium root rot in squash using mycorrhizal fungi and antagonistic microorganisms", Egyptian Journal of Biological Pest Control, 2022  
Publication 1 %
- 
- 6** Yosep Seran Mau, Antonius Simon Sebho Ndiwa, Jenny Ema Rut Markus, Shirly Seahan Oematan et al. "Blast Resistance Levels of Red and Black Upland Rice Local Cultivars from Indonesia", Asian Journal of Crop Science, 2018  
Publication <1 %
- 
- 7** "Rice Production Worldwide", Springer Science and Business Media LLC, 2017  
Publication <1 %
- 
- 8** Salma Javed, Tabassum Ara Khanum, Ashraf Ali. "Storage and efficacy of entomopathogenic nematode species as a biocontrol agent against the armyworm, Spodoptera litura (FABRICIUS) (Lepidoptera: Noctuidae)", Egyptian Journal of Biological Pest Control, 2022  
Publication <1 %
-

9

Cuong Van Pham, Seiichi Murayama, Yukio Ishimine, Yoshinobu Kawamitsu, Keiji Motomura, Eiji Tsuzuki. " Sterility of Thermo-Sensitive Genic Male Sterile Line, Heterosis for Grain Yield and Related Characters in F Hybrid Rice ( L.) ", Plant Production Science, 2015

Publication

<1 %

10

Esmat Hegazy, Wedad Khafagi, Essam Agamy. "Arthropod fauna of Thymelaea hirsuta in the Egyptian western desert, with a special reference to *Olpium kochi*", Egyptian Journal of Biological Pest Control, 2022

Publication

<1 %

11

"Trichoderma: Agricultural Applications and Beyond", Springer Science and Business Media LLC, 2020

Publication

<1 %

12

Babak Nakhoda, Hei Leung, Merlyn S. Mendioro, Ghasem Mohammadi-nejad, Abdelbagi M. Ismail. "Isolation, characterization, and field evaluation of rice (*Oryza sativa* L., Var. IR64) mutants with altered responses to salt stress", Field Crops Research, 2012

Publication

<1 %

13

C. Ndikuryayo, M. Ochwo-Ssemakula, P. Gibson, J. Lamo. "Resistance to Rice yellow mottle virus and performance of selected

<1 %

improved rice genotypes in central Uganda",  
Crop Protection, 2020

Publication

---

14

Microbial Diversity and Biotechnology in Food Security, 2014.

Publication

---

<1 %

15

Mohamed T. El-Saadony, Ahmed M. Saad, Soliman M. Soliman, Heba M. Salem et al. "Plant growth-promoting microorganisms as biocontrol agents of plant diseases: Mechanisms, challenges and future perspectives", Frontiers in Plant Science, 2022

Publication

---

<1 %

16

Moutoshi Chakraborty, S. M. Fajle Rabby, Dipali Rani Gupta, Mahfuzur Rahman et al. "Natural Protein Kinase Inhibitors, Staurosporine, and Chelerythrine Suppress Wheat Blast Disease Caused by Magnaporthe oryzae Triticum", Microorganisms, 2022

Publication

---

<1 %

17

M. Ni, K. Gu, B. Hassan, D. Ning, Y. Zheng, Y. Qi, Y. Xu. "Effect of oviposition by Bactrocera dorsalis on the antioxidant activity of orange juice", Brazilian Journal of Biology, 2019

Publication

---

<1 %

18

Yosep Seran Mau, Antonius Simon Sebh. "Field Evaluation of Late Leaf Spot and Leaf Rust Resistance and the Associated Yield

<1 %



Losses in Indonesian Groundnut Genotypes",  
Asian Journal of Plant Sciences, 2018

Publication

---

19

Afroz Rais, Muhammad Shakeel, Fauzia Yusuf Hafeez, Muhammad Nadeem Hassan. "Plant growth promoting rhizobacteria suppress blast disease caused by *Pyricularia oryzae* and increase grain yield of rice", BioControl, 2016

Publication

---

20

Asep K. Supriatna, Ema Carnia, Meksianis Z. Ndi. "Fibonacci numbers: A population dynamics perspective", Heliyon, 2019

Publication

---

21

Somayeh Dariush, Mostafa Darvishnia, Ali-Akbar Ebadi, Fereidoun Padasht-Dehkaei, Eidi Bazgir. "Screening brown spot resistance in rice genotypes at the seedling stage under water stress and irrigated conditions", Archives of Phytopathology and Plant Protection, 2020

Publication

---

22

Benish Ashfaq, Hafiz M. Imran Arshad, M. Atiq, Sumaira Yousaf, Kamran Saleem, Amna Arshad. "Biochemical Profiling of Resistant Phenotypes Against *Bipolaris oryzae* Causing Brown Spot Disease in Rice", Frontiers in Agronomy, 2021

Publication

<1 %

<1 %

<1 %

<1 %

---

23 A. Moneim Babu Fatih. "Genotypic stability analysis of yield and related agronomic characters in wheat-Agropyron derivatives under varying watering regimes", Theoretical and Applied Genetics, 1987 <1 %  
Publication

---

24 Md. Motaher Hossain. "Systemic resistance to bacterial leaf speck pathogen in Arabidopsis thaliana induced by the culture filtrate of a plant growth-promoting fungus (PGPF) Phoma sp. GS8-1", Journal of General Plant Pathology, 06/2008 <1 %  
Publication

---

25 Muhammad Imran, Kamal A. M. Abo-Elyousr, Magdi A. Mousa, Maged M. Saad. "Screening and biocontrol evaluation of indigenous native Trichoderma spp. against early blight disease and their field assessment to alleviate natural infection", Egyptian Journal of Biological Pest Control, 2022 <1 %  
Publication

---

26 P. Plumb - Dhindsa, A. M. Mondjane. "Index of plant diseases and associated organisms of Mozambique", Tropical Pest Management, 2008 <1 %  
Publication

---

27 Gamil A. Amin, Narmen A. Youssef, Saleh Bazaid, Waleed D. Saleh. "Assessment of <1 %

insecticidal activity of red pigment produced by the fungus *Beauveria bassiana*", *World Journal of Microbiology and Biotechnology*, 2010

Publication

---

28

Perera, UIP, Disna Ratnasekera, SGJN Senanayake, and D Sumith de Z Abeysiriwardena. "Genetic parameters and correlations of yield attributing characteristics of weedy rice in Sri Lanka", *Journal of the National Science Foundation of Sri Lanka*, 2014.

Publication

---

29

N Siswanto, N Bintoro, JNW Karyadi, S Rahayoe, Suryanti. "The kinetics of pesticides and UVC irradiation effect on the rate of shallot (*Allium cepa* var *ascalonicum* L. Back) seed quality change during storage", *IOP Conference Series: Earth and Environmental Science*, 2021

Publication

---

30

Truong, Hoai, and Evelyn Gergon. "Current Research on Fungal Pathogens Associated with Rice", *Fungi From Different Substrates*, 2014.

Publication

---

<1 %

<1 %

<1 %

---

Exclude quotes      On

Exclude matches      < 10 words

Exclude bibliography      On